



831172 001N

# A Subroutine Package for the Efficient Solution of the Eigenproblem of Real Symmetric Toeplitz Matrices

Marvin J. Goldstein

William A. Babson

Computer and Information Services Department

REFERENCE ONLY



**Naval Underwater Systems Center**  
Newport, Rhode Island/New London, Connecticut

<b>Report Documentation Page</b>			Form Approved OMB No. 0704-0188	
<p>Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p>				
1. REPORT DATE <b>30 DEC 1983</b>	2. REPORT TYPE <b>Technical Memo</b>	3. DATES COVERED <b>30-12-1983 to 30-12-1983</b>		
4. TITLE AND SUBTITLE <b>A Subroutine Package for the Efficient Solution of the Eigenproblem of Real Symmetric Toeplitz Matrices</b>			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) <b>Marvin Goldstein; William Babson</b>			5d. PROJECT NUMBER <b>771Y00</b>	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Underwater Systems Center, New London, CT, 06320</b>			8. PERFORMING ORGANIZATION REPORT NUMBER <b>TM 831172</b>	
			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
			12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>	
13. SUPPLEMENTARY NOTES <b>NUWC2015</b>				
14. ABSTRACT <b>We discuss a subroutine package for solving the eigenproblem of bisymmetric Toeplitz matrices. When the eigenvalues and eigenvectors of a bisymmetric Toeplitz matrix are computed, storage requirements and execution time are reduced by taking advantage of the special structure of the matrix.</b>				
15. SUBJECT TERMS <b>bisymmetric Toeplitz matrices</b>				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>16</b>
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	19a. NAME OF RESPONSIBLE PERSON	

NAVAL UNDERWATER SYSTEMS CENTER  
NEW LONDON LABORATORY  
NEW LONDON, CONNECTICUT 06320

Technical Memorandum

A SUBROUTINE PACKAGE FOR THE EFFICIENT SOLUTION OF  
THE EIGENPROBLEM OF REAL SYMMETRIC TOEPLITZ MATRICES

Date: December 30, 1983

Prepared by: Marvin J. Goldstein  
Marvin J. Goldstein

William A. Babson Jr.  
William A. Babson

Approved for public release; distribution unlimited

ABSTRACT

We discuss a subroutine package for solving the eigenproblem of bisymmetric Toeplitz matrices. When the eigenvalues and eigenvectors of a bisymmetric Toeplitz matrix are computed storage requirements and execution time are reduced by taking advantage of the special structure of the matrix.

ADMINISTRATIVE INFORMATION

This memorandum was prepared under Job Order No. 771Y00, Special Projects and Studies. The authors are located at the Naval Underwater Systems Center, New London, Connecticut, 06320.

ACKNOWLEDGEMENT

The authors are grateful to Dr. John Ianniello (Code 3212) for technically reviewing this memorandum.

## INTRODUCTION

An  $n$ -th order real symmetric matrix  $M = (M_{jk})$  for which

$$M_{jk} = M_{j+1, k+1} \quad (1)$$

is called a bisymmetric Toeplitz matrix. Bisymmetric Toeplitz matrices arise, for example, as correlation matrices in the study of discrete stationary random processes. Each element of the correlation matrix is a function of  $j-k$ ; that is,

$$M_{jk} = M_{j-k} = M_{k-j}$$

(e.g.  $M_{jk} = \cos(j-k)c$ , where  $c$  is a real number.)

In [1], using the bisymmetry of matrix  $M$ , Goldstein shows that the eigenvalues and eigenvectors of matrix  $M$  can be computed from two real symmetric matrices one-half its size, if it is of even order. In this memorandum, we provide a FORTRAN subroutine package that is an implementaiton of this reduction. By using this package, one reduces both computer time and storage ordinarily needed to calculate the eigenvalues and eigenvectors of bisymmetric Toeplitz matrices.

## ALGORITHM

The sub-program package in the Appendix computes the eigenvalues and (optionally) the eigenvectors of an even order,  $n$ , real symmetric matrix  $M$ , whose coefficients satisfy Eq. (1), from two real symmetric matrices:

$$M_{jk}^{\sim}(+) = M_{jk} + M_{j,n+1-k} = M_{kj}^{\sim}(+) \quad (2)$$

( $k=1, \dots, n/2; j=1, \dots, k$ )

$$M_{jk}^{\sim}(-) = M_{jk} - M_{j,n+1-k} = M_{kj}^{\sim}(-) \quad (3)$$

one-half the size of matrix  $M$ , by [1, 2].

The matrices  $M^{\sim}(+)$  and  $M^{\sim}(-)$  are computed from the first row elements of matrix  $M$ . For example, by Property (1), Formula (2) reduces to

$$M_{jk}^{\sim}(+) = M_{1,k-(j-1)} + M_{1,n-(k+j)+2} \quad (4)$$

$$= M_{kj}^{\sim}(+) \quad k=1, \dots, n/2; j=1, \dots, k$$

where addition of the coefficients of the first row of matrix  $M$  is replaced by subtraction when computing  $M^{\sim}(-)$ .

Forming  $M\sim(+)$  first, the subroutine package calculates the eigenvalues and (optionally) the eigenvectors of  $M\sim(+)$ . Then  $M\sim(-)$  is formed and its eigenvalues and (optionally) eigenvectors are computed.

The eigenvalues and eigenvectors of matrices  $M\sim(+)$  and  $M\sim(-)$  determine the eigenvalues and eigenvectors of matrix  $M$ . If  $z$  is an eigenvector of  $M\sim(+)$  with corresponding eigenvalue  $d$ , and  $J$  is the identity matrix of order  $n/2$  with its columns written in reverse order, then

$$\begin{aligned} & \begin{bmatrix} z \\ Jz \end{bmatrix} = d \begin{bmatrix} z \\ Jz \end{bmatrix}. \\ M \begin{bmatrix} z \\ Jz \end{bmatrix} &= d \begin{bmatrix} z \\ Jz \end{bmatrix}. \end{aligned} \tag{5}$$

Similarly, if  $z$  is an eigenvector of  $M\sim(-)$  with corresponding eigenvalue  $d$ , then

$$\begin{aligned} & \begin{bmatrix} z \\ -Jz \end{bmatrix} = d \begin{bmatrix} z \\ -Jz \end{bmatrix}. \\ M \begin{bmatrix} z \\ -Jz \end{bmatrix} &= d \begin{bmatrix} z \\ -Jz \end{bmatrix}. \end{aligned} \tag{6}$$

## STORAGE CONSIDERATIONS

Note that in (5) and (6), since premultiplying a vector by matrix  $J$  just reverses the order of the vector's components, all the information about the eigenvectors of matrix  $M$  is given by the eigenvectors of matrices  $M\sim(+)$  and  $M\sim(-)$ ; therefore, only  $n/2$  of the components of each eigenvector of matrix  $M$  have to be stored, and a fortiori, only  $n^{**}2/2$  entries (instead of  $n^{**}2$ ) are required to store all the eigenvectors of matrix  $M$ .

Furthermore, since the eigenvalues and eigenvectors of  $M\sim(+)$  are computed independently of those for  $M\sim(-)$ , matrices  $M\sim(+)$  and  $M\sim(-)$  may share the same storage area for a matrix of order  $n/2$ . Therefore, the reduction requires a maximum of  $(3/4)n^{**}2 + n$  storage locations to accomodate the matrices  $M\sim(+)$ ,  $M\sim(-)$  and their eigenvalues and eigenvectors, instead of the  $2n^{**}2 + n$  required when the eigensystem is solved directly from matrix  $M$ . In particular, only  $(1/4)n^{**}2 + n$  storage locations are used to accomodate  $M\sim(+)$  and  $M\sim(-)$  and their eigenvalues, if only the eigenvalues of matrix  $M$  are required.

## TIMING

Computational results indicate that the eigenproblem for a large matrix  $M$  can be solved by computer at least four times faster this way than directly from matrix  $M$ . For example, the following table gives representative execution times (in seconds) to compute the eigenvalues of bisymmetric Toeplitz matrices of orders 16, 32, 64, 128, 256 by the reduction SUBROUTINE

MATRED (in the Appendix) and directly by the IMSL library SUBROUTINE EIGRS [3] on the VAX 11/780 in double precision arithmetic.

Order (n)	MATRED (using EIGRS)	EIGRS
16	.03	.07
32	.13	.38
64	.73	2.64
128	4.97	21.00
256	38.53	248.63

The following table gives representative execution times (in seconds) for a single precision version of SUBROUTINE MATRED and a single precision version of EIGRS on the UNIVAC 1100/62:<sup>\*</sup>

Order (n)	Single Precision MATRED	Single Precision EIGRS
16	.02	.05
32	.10	.33
64	.60	2.24
128	4.08	16.76
256	30.16	129.04

---

\* Since only a single precision version of EIGRS is available in the IMSL library on the UNIVAC, the UNIVAC version of MATRED is single precision.

In multi-program computer environments, variations in the execution time of a program can occur due to differences in machine load. Therefore, in order to smooth out these variations, subroutines MATRED and EIGRS were run a number of times on different matrices of order n and the average execution time for each order was recorded in the tables.

#### USER PROCEDURE

The calling sequence of SUBROUTINE MATRED (double precision version) is defined in its commentary in the Appendix. The relocatable code, MATRD.OBJ, is in the VAX 11/780 directory [MJG] on node 2. The UNIVAC single precision version of SUBROUTINE MATRED is in the UNIVAC 1100/62 (node 2) file MGOLDSTN\*MATRD and has the element name MATRD. It's calling sequence is the same as the double precision version on the VAX, but none of the arguments in the sequence are double precision.

#### SUMMARY

A subroutine package for computing the eigenvalues and (optionally) the eigenvectors of a bisymmetric Toeplitz matrix is available which reduces storage requirements and execution time by taking advantage of the special structure of the matrix. In particular, if only the eigenvalues of the matrix are required storage requirements for arrays and execution time are reduced by 75% when the matrix is sufficiently large.

REFERENCES

1. Goldstein Marvin, "Reduction of the Eigenproblem for Hermitian Persymmetric Matrices," *Math, Comp.*, v. 38, Jan. 1974
2. Goldstein Marvin, "Further Decomposition of the Pseudoinverse and Eigensystem of a Hermitian Persymmetric Matrix," *ACM SIGNUM Newsletter*, v. 13, no. 1, March 1978.
3. "EIGRS," *IMSL Library for VAX-11/780*, Eighth Edition, 1980.

## APPENDIX

```

100      SUBROUTINE MATRED(IVEC,MR1,N,IFULST,MTILFL,IFUL,MTILSY,D,Z,IZ,WK)
200      C ****
300      C      A SUBROUTINE PACKAGE FOR THE EFFICIENT SOLUTION
400      C      OF THE EIGENPROBLEM OF REAL SYMMETRIC TOEPLITZ MATRICES
500      C
600      C      BY
700      C
800      C      MARVIN GOLDSTEIN AND WILLIAM BABSON
900      C
1000     C *** THIS SUB-PROGRAM PACKAGE COMPUTES THE EIGENVALUES AND      ***
1100     C *** (OPTIONALLY) THE EIGENVECTORS OF AN EVEN ORDER (n) REAL SYM-  ***
1200     C *** METRIC TOEPLITZ MATRIX:                                     ***
1300     C
1400     C      M = M                                         (1)
1500     C      jk   j+1, k+1
1600     C
1700     C      = M
1800     C      k j
1900     C
2000     C *** FROM TWO REAL SYMMETRIC MATRICES:                      ***
2100     C
2200     C      M~ (+) = M + M = M~ (+)                                (2)
2300     C      jk       jk   j,n+1-k   kj
2400     C
2500     C      (k=1, ..., n/2; j=1, ..., k)
2600     C
2700     C      M~ (-) = M - M = M~ (-)                                (3)
2800     C      jk       jk   j,n+1-k   kj
290     C
310     C *** ONE-HALF THE SIZE OF MATRIX M, BY [1, 2].           ***
3200     C
3300     C *** THE MATRICES M~ (+) AND M~ (-) CAN BE COMPUTED FROM THE  ***
3400     C *** FIRST ROW ELEMENTS OF MATRIX M. FOR EXAMPLE, BY PROPERTY (1), ***
3500     C *** FORMULA (2) REDUCES TO
3600     C      M~ (+) = M + M                                         (4)
3700     C      jk   1,k-(j-1)   1,n-(k+j)+2
3800     C
3900     C      = M~ (+)          k=1, ..., n/2; j=1, ..., k
4000     C      k j
4100     C
4200     C *** WHERE ADDITION OF THE COEFFICIENTS OF THE FIRST ROW OF MATRIX ***
4300     C *** M IS REPLACED BY SUBTRACTION WHEN COMPUTING M~ (-).        ***
4400     C
4500     C *** IF z IS AN EIGENVECTOR OF M~ (+) WITH CORRESPONDING EIGEN- ***
4600     C *** VALUE d, AND J IS THE IDENTITY MATRIX OF ORDER n/2 WITH ITS  ***
4700     C *** COLUMNS WRITTEN IN REVERSE ORDER, THEN
4800     C
4900     C      | z |   | z |
5000     C      M |   | = d |   |.                               (5)
5100     C      | Jz |   | Jz |
5200     C
5300     C *** SIMILARLY, IF z IS AN EIGENVECTOR OF M~ (-) WITH CORRESPONDING ***
5400     C *** EIGENVALUE d, THEN
5500     C
5600     C      | z |   | z |
5700     C      M |   | = d |   |.                               (6)
5800     C      | -Jz |   | -Jz |
5900     C
6000     C *** COMPUTATIONAL RESULTS SHOW THAT THE EIGENPROBLEM FOR      ***
6100     C *** A MATRIX THAT SATISFIES (1) CAN BE SOLVED BY COMPUTER AT LEAST ***
6200     C *** FOUR TIMES FASTER THIS WAY THAN BY SOLVING THE EIGENPROBLEM    ***
6300     C *** DIRECTLY FROM M FOR LARGE VALUES OF n.                         ***
6400     C

```



```

14300 C
14400 C      INTEGER N,IER,IFULST,IFUL,IZ,IVEC,NDIV2,IPOINT
14500 C      DOUBLE PRECISION MR1,MTILFL,MTILSY,D,Z,WK,SIGN
14600 C      DIMENSION MR1(1),MTILFL(IFUL,1),MTILSY(1),D(1),Z(IZ,1),WK(1)
14700 C
14800 C      COMPUTE ORDER OF REDUCED MATRICES
14900 C
15000 C      NDIV2=N/2
15100 C
15200 C      SET SIGN TO COMPUTE M~(+)
15300 C
15400 C      SIGN=1.0 DO
15500 C
15600 C      FIRST TIME THRU LOOP SOLVE EIGEN-
15700 C      PROBLEM FOR M-(+); SECOND TIME FOR
15800 C      M~(-)
15900 C
16000 DO 40 I=1,2
16100   IPOINT=(I-1)*NDIV2+1
16200   IF(IFULST.LT.1)GO TO 10
16300   GO TO 20
16400 C      THEN COMPUTE USING SYMMETRIC STORAGE MODE
16500   10    CALL MTLSYM(NDIV2,MR1,MTILSY,SIGN)
16600   CALL EIGRS(MTILSY,NDIV2,IVEC,D(IPOINT),Z(1,IPOINT),IZ,WK,IER
16700   +
16800   GO TO 30
16900 C      ELSE COMPUTE USING FULL STORAGE MODE
17000   20    IVEC=IVEC+10
17100 C
17200 C      SINCE IMSL ROUTINE EIGRS REQUIRES
17300 C      THAT MTILFL BE EXACTLY N/2 BY N/2
17400 C      THE FOURTH ARGUMENT IN THE MTLFL
17500 C      CALL HAS BEEN REPLACED BY NDIV2.
17600 C      RESTORE THIS ARGUMENT TO IFUL IF
17700 C      REQUIRED BY AN EIGENVALUE ROUTINE.
17800 C
17900   CALL MTLFUL(NDIV2,MR1,MTILFL,NDIV2,SIGN)
18000   CALL EIGRS(MTILFL,NDIV2,IVEC,D(IPOINT),Z(1,IPOINT),IZ,WK,IER
18100   +
18200   30    CONTINUE
18300 C
18400 C      SET SIGN TO COMPUTE M~(-)
18500 C
18600 C      SIGN=-1.0 DO
18700 C
18800 C      40 CONTINUE
18900 C
19000 C      RETURN
19100 C      END

```

```

19200      SUBROUTINE MTLSYM(NDIV2,MR1,MTILDA,SIGN)
19300      C **** THIS SUBROUTINE RETURNS M~(+) OR M~(-) STORED IN ***
19400      C ***SYMMETRIC STORAGE MODE IN THE ONE-DIMENSIONAL, DOUBLE ***
19500      C ***PRECISION ARRAY MTILDA. M~(+) AND M~(-) ARE COMPUTED ***
19600      C ***FROM THE FIRST ROW OF MATRIX M, NAMELY MR1, WHEN SIGN ***
19700      C ***IS +1 AND -1, RESPECTIVELY.
19800      C ****
19900      C ****
20000      INTEGER I,J,K,NDIV2,N,L1,L2
20100      DOUBLE PRECISION MR1,MTILDA,SIGN
20200      DIMENSION MR1(1),MTILDA(1)
20300      C
20400      N=NDIV2*2
20500      C
20600      I=0
20700      DO 20 K=1,NDIV2
20800      DO 10 J=1,K
20900      I=I+1
21000      L1=K-J+1
21100      L2=N+2-K-J
21200      MTILDA(I)=MR1(L1)+MR1(L2)*SIGN
21300      10    CONTINUE
21400      20    CONTINUE
21500      C
21600      RETURN
21700      END

21800      SUBROUTINE MTLFUL(NDIV2,MR1,MTILDA,IFUL,SIGN)
21900      C **** THIS SUBROUTINE RETURNS M~(+) OR M~(-) STORED IN ***
22000      C ***FULL STORAGE MODE IN THE TWO-DIMENSIONAL, DOUBLE PRE- ***
22100      C ***CISION ARRAY MTILDA. M~(+) OR M~(-) ARE COMPUTED FROM ***
22200      C ***THE FIRST ROW OF MATRIX M, NAMELY MR1, WHEN SIGN IS ***
22300      C ***+1 AND -1, RESPECTIVELY.
22400      C ****
22500      C ****
22600      C
22700      INTEGER N,NDIV2,I,IFUL,J,K,L1,L2
22800      DOUBLE PRECISION MR1,MTILDA,SI,N
22900      DIMENSION MR1(1),MTILDA(IFUL,1)
23000      C
23100      N=NDIV2*2
23200      C
23300      DO 20 K=1,NDIV2
23400      DO 10 J=1,K
23500      L1=K-J+1
23600      L2=N+2-K-J
23700      MTILDA(J,K)=MR1(L1)+MR1(L2)*SIGN
23800      MTILDA(K,J)=MTILDA(J,K)
23900      10    CONTINUE
24000      20    CONTINUE
24100      C
24200      RETURN
24300      END

```

A SUBROUTINE PACKAGE FOR THE EFFICIENT SOLUTION OF  
THE EIGENPROBLEM OF REAL SYMMETRIC TOEPLITZ MATRICES  
Marvin J. Goldstein, Computer and Information Services Dept.  
William Babson, Computer and Information Services Dept.  
TM No. 831172  
UNCLASSIFIED

DISTRIBUTION LIST

External:

Mr. I. L. Avrunin  
David Taylor Naval Ship  
Research & Development  
Center  
Headquarters  
Bethesda, MD 20084

Dr. Martin Schultz  
Research Center for Scientific Comp.  
Yale Univ.  
P. O. Box 2158 Yale Station  
New Haven, CT 06520

Mr. Ken Kunec  
Naval Data Automation  
Command  
Washington Navy Yard  
Washington, DC 20374

Internal:

Code 00	CAPT Ailes	Code 33	J.E. Miller	60	Dr. D. Bordelon
01	E.L. Messere		Dr. H.F. Dwyer		Dr. Jeffrey Cohen
01A	John Keegan		Dr. C. Carter	70	Cort R. DeVoe
039	Dr. D.M. Viccone		A. Quazi	701	R. Wilson Npt
10	Dr. W.A. VonWinkle		Peter Stahl		R. Streit
101	Dr. E. Eby		J. Wolcin		D. McCue
02	D.C. Kindilien		Dr. M.B. Moffett		J. Babiec Npt
0211	R.H. Bernier		C.J. Becker	71	G. Elias Npt
02	NL Library (3 copies)		F.C. Gannon		M. Goldstein
021311	NPT Library		R.J. MacDonald		(15 Copies)
			J. Saikowski		
20	W.L. Clearwaters		G. Botseas		D. Drinkard
234	R.J. Conley		Dr. D. Lee		D. Aker
32	A. Lesick		J. Nordquist		
	Dr. N.L. Owsley		E.R. Robinson		N. Sulinski
	Dr. John Ianniello		H. Weinberg		W. Babson (15 Copies)
	J. K. Sullivan		H. Sternberg		E. Montavon
	H. Axtell	34	Dr. David Wood		J. Sikorski
	R.C. Cox		Michael Fecher		Louis Ledoux Npt
	James Pearson		J.R. Katan		R. Johnson
	L.C. Ng		Dr. P.E. Fessenden		H. Eley
	J.V. Sanchis		C.N. Flemings		Cynthia Borges Npt
	B. Helme, Jr.		K.F. Hafner		Marcel Nadeau Npt
	Dr. C.H. Sherman		John Casey		John Ventura Npt
	W. Fox	35	Anthony Bruno		Phoebe Liu Npt
	B.G. Buehler		D. Dixon		John Sabulis Npt
	A.W. Ellenthalpe	36	Tom Conrad	72	Gordon Daglieri Npt
	Dr. Peter G. Cable		L. Cabral		S. Schneller Npt
	Dr. S. Ko		R.P. Eidimtas		J. Auwood
	D.T. Porter		H.A. Rosen		T. Wheeler
	W.H. Wharton		R.V. Cherry		R. Warren Npt
	T. Anderson		W.J. Ryan		P. Breslin Npt
	J. Ionata		J. Griffin		A. Alfiero
	M. Kuuznitz		D.G. Blundell		R. Clark
	J. Gannon		M.P. Lydon		J. Gribbin
	C. Bowman		S.I. Wax		D. Quigley
	P.R. Miner		S.E. Ashton	73	S. Capizzano
	Rosemary Molino	37	Raymond McMahon		J. MacDonald
	Jose Munoz		S. Meyers		R. Pingree
	J. Shores	38	C.M. Curtis		R. Cote
	R. Leask	401	Q. Huynh		R. Forget
	Russell Christman		J.E. Sims		A. Sullivan
33	Dr. A.H. Nuttal		A.D. Carlson		J. Surdo
	J.B. Paniszczyn		J.H. Clark		M. Lee
	Dr. F.R. DiNapoli		Dr. A.J. Kalinowski	74	C. Brockway
	E.P. Jensen		Dr. R.G. Kasper		M. Becker
	E.G. Kanabis		R.R. Manstan		L. Doyle
	S.C. Gerengher		E.L. McLaughlin		J. Koonce
	J. Gregor		R.S. Munn		J. Pappadia
	W.A. Goldman		C.W. Nebelung		A. Levander
	P.M. Anchors		Dr. J.S. Patel		Norman Dube Npt
	J.J. Higgs		B.A. Radley		A. Blau Npt
	I.B. Cohen	402	A.Y. Shigematsu		D. Stanhope Npt
	D.W. Counsellor	4111	M.A. Tucchio		R. Hoy Npt
	Dr. R. Radliniski	4331	M. Berger		N. Bradbury Npt
	D.A. Fingerman		Samuel Horvitz		
	R.J. Trembley	5202	B.L. Antrim		
			S.B. Walsh		
			Stephan Schady		